

The need for clean air: The way air pollution and climate change affect allergic rhinitis and asthma

Ibon Eguiluz-Gracia¹ | Alexander G. Mathioudakis^{2,3} | Sabine Bartel^{4,5} |
Susanne J. H. Vijverberg⁶ | Elaine Fuertes⁷ | Pasquale Comberiati^{8,9} |
Yutong Samuel Cai^{10,11} | Peter Valentin Tomazic¹² | Zuzana Diamant^{13,14} |
Jørgen Vestbo^{2,3} | Carmen Galan¹⁵ | Barbara Hoffmann¹⁶

¹Allergy Unit, IBIMA-Hospital Regional Universitario de Malaga-UMA, Malaga, Spain

²Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, The University of Manchester, Manchester Academic Health Science Centre, UK

³North West Lung Centre, Wythenshawe Hospital, Manchester University NHS Foundation Trust, Southmoor Road, Manchester, UK

⁴Early Life Origins of Chronic Lung Disease, Research Center Borstel, Leibniz Lung Center, Member of the German Research Center for Lung Research (DZL), Borstel, Germany

⁵Department of Pathology and Medical Biology, University Medical Center Groningen, GRIAC Research Institute, University of Groningen, Groningen, The Netherlands

⁶Department of Respiratory Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

⁷National Heart and Lung Institute, Imperial College London, London, UK

⁸Section of Paediatrics, Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy

⁹Department of Clinical Immunology and Allergology, Sechenov University, Moscow, Russia

¹⁰Department of Epidemiology and Biostatistics, MRC Centre for Environment and Health, School of Public Health, Imperial College London, London, UK

¹¹The George Institute for Global Health, University of Oxford, Oxford, UK

¹²Department of General ORL, Head and Neck Surgery, Medical University of Graz, Graz, Austria

¹³Department of Respiratory Medicine & Allergology, Institute for Clinical Science, Skane University Hospital, Lund University, Lund, Sweden

¹⁴Department of Respiratory Medicine, First Faculty of Medicine, Charles University and Thomayer Hospital, Prague, Czech Republic

¹⁵Department of Botany, Ecology and Plant Physiology, International Campus of Excellence on Agrifood (ceiA3), University of Córdoba, Córdoba, Spain

¹⁶Institute for Occupational, Social and Environmental Medicine, Medical Faculty, University of Düsseldorf, Düsseldorf, Germany

Correspondence

Ibon Eguiluz-Gracia, Allergy Unit, IBIMA-Hospital Regional Universitario de Malaga-UMA, ARADyAL, Malaga, Spain.
Email: iboneguiluz@gmail.com

Alexander G. Mathioudakis, Division of Infection, Manchester Academic Health Science Centre, Immunity and Respiratory Medicine, School of Biological Sciences, Manchester University NHS Foundation Trust, University of Manchester, Manchester, UK.
Email: Alexander.Mathioudakis@Manchester.ac.uk

Abstract

Air pollution and climate change have a significant impact on human health and well-being and contribute to the onset and aggravation of allergic rhinitis and asthma among other chronic respiratory diseases. In Westernized countries, households have experienced a process of increasing insulation and individuals tend to spend most of their time indoors. These sequelae implicate a high exposure to indoor allergens (house dust mites, pets, molds, etc), tobacco smoke, and other pollutants, which have an impact on respiratory health. Outdoor air pollution derived from traffic and other human activities not only has a direct negative effect on human health but also enhances the allergenicity of some plants and contributes to global warming.

Eguiluz-Gracia and Mathioudakis are equally contributed to this work.

This review was produced as result of the collaboration between the European Respiratory Society (ERS) Early Career Members Committee (ERS ECMC) and the European Academy of Allergy and Clinical Immunology (EAACI) Junior Members Assembly (EAACI JMA) Board, as part of an Environmental Awareness Initiative. It is not an official document of the ERS or EAACI and the views expressed are those of the authors and not necessarily those of the ERS or EAACI.

Funding information

Instituto de Salud Carlos III; Spanish Ministry of Science and Innovation through the Rio Hortega; RETICS schemes, Grant/Award Number: CM17/00140 and RD16/0006/0001; National Institute of Health Research Manchester Biomedical Research Centre; Medical Research Council Early Career Research Fellowship awarded through the MRC-PHE Centre for Environment and Health, Grant/Award Number: MR/M501669/1

Climate change modifies the availability and distribution of plant- and fungal-derived allergens and increases the frequency of extreme climate events. This review summarizes the effects of indoor air pollution, outdoor air pollution, and subsequent climate change on asthma and allergic rhinitis in children and adults and addresses the policy adjustments and lifestyle changes required to mitigate their deleterious effects.

KEYWORDS

allergic rhinitis, asthma, climate change., environment, pollution

1 | INTRODUCTION

Since the beginning of the industrial revolution, Western countries experienced an explosive process of urbanization, which dramatically affected environmental exposures. Following this trend, many low-to-middle income countries are undergoing similar processes. Consequently, >90% of the population lives in places where air quality does not meet the recommendations of the *World Health Organization* (WHO).¹ The *European Environmental Agency* reported that most urban dwellers were exposed to concentrations of fine particulate matter (PM_{2.5}) and particulate matter of ≤10 mm in diameter (PM₁₀) above WHO recommendations (74% and 42%, respectively).² Importantly, air pollution is currently one of the leading causes of premature death in the world.^{3,4}

Allergic rhinitis (AR) and asthma share many pathophysiological links^{5,6} and are among the commonest respiratory conditions,^{7,8} with their increasing prevalence mirroring the rise in Westernized lifestyle worldwide.⁹ Because the airways represent one of the major boundaries of the body, environmental exposures (collectively termed “the exposome”)¹⁰ greatly affect the homeostasis of the respiratory mucosae. Importantly, climate and urban dwelling (with its associated decrease in biodiversity) significantly determine the exposome composition.¹¹ Among the exposome components, pollutants, microbes, and allergens have a substantial impact on health.¹²

Several policy changes could help reduce the deleterious components of the exposome and minimize their effects on respiratory health.² Of note, some policy measures have already proven effective at decreasing the burden of air pollution-related diseases (e.g., restrictions on tobacco smoking in public places).¹³

This narrative review summarizes the latest insights regarding the effects of indoor and outdoor pollution and climate change on AR and asthma, and addresses the policy adjustments required to mitigate their effects. To this end, we identified relevant articles published during the period 2014-2019, together with several previous key studies related to the topic.

2 | INDOOR AIR POLLUTION

Most individuals in Westernized countries spend ~80% of their time indoors,^{14,15} demonstrating the importance of indoor air quality.

The composition of indoor air is affected by several factors including outdoor pollutants, the quality/quantity of ventilation, indoor allergens, and activities such as smoking, heating, and cooking.¹⁶

2.1 | Second-hand exposure to tobacco smoke**2.1.1 | Epidemiological evidence**

Tobacco smoke contains at least 4500 toxic chemical compounds, including PM, oxidative gases, heavy metals, and at least 50 carcinogens.¹⁷ Tobacco smoke poses significant health risks to nonsmokers who inhale the smoke in various microenvironments, such as households or workplaces (second-hand smoke (SHS) exposure). Recently, the pyro-synthesis and cigarette combustion related to domestic smoking were identified as key phenomena increasing the levels of PM and toxic chemical agents in households.¹⁸

Second-hand smoke exposure during pregnancy and infancy is associated with asthma onset, poor asthma control, and more severe exacerbations during childhood,¹⁹⁻²¹ among other chronic conditions.²² Prenatal and postnatal SHS exposure was linked to a 21%-85% increase in the risk of asthma in children, with the highest effect observed among children exposed to tobacco smoke during the first two years of life.^{20,23} More recently, a study of five European birth cohorts (n = 10 860) showed that maternal smoking during infancy correlated with a 15% (95%CI: 0%-31%) increase in the risk of asthma in children.²⁴ Interestingly, SHS exposure might induce epigenetic changes with transgenerational repercussions on asthma onset,²⁵ which would imply a very long-term effect of tobacco smoking on respiratory health. The immaturity of the immune and respiratory systems of children and their larger air volume per weight kilogram inhaled as compared to adults might explain the high sensitivity to tobacco smoke during childhood.²⁶ The initial studies conducted by the *European Community Respiratory Health Survey* (ECRHS) in different European countries did not reach a definitive conclusion regarding the role of SHS exposure on asthma in adults.²⁷ Conversely, later studies suggested a higher risk of adult-onset asthma in patients exposed to tobacco smoke,^{28,29} together with a slightly lower risk of seasonal AR.²⁹ Similarly, a recent study from the ECRHS found associations between SHS exposure and physician-diagnosed asthma and poorer asthma control in adults, yet no effect was observed on the lung function.³⁰

2.1.2 | Tobacco smoke alters airway microbiota

Similar to the gut,^{31,32} the respiratory tract of healthy humans is colonized by a variety of different bacteria, viruses, and fungi.³³ This colonization may shift in response to changes in the local environment (e.g., tobacco smoke), potentially inducing a long-lasting state of bacterial dysbiosis.³⁴ A study reported that the pharyngeal microbiota of individuals exposed to tobacco smoke was richer in species, such as *Porphyromonas*, *Neisseria*, and *Gemella*, compared to nonexposed subjects, but the authors failed to identify significant changes in the microbiota recovered from bronchoalveolar lavage samples.³⁵ Various tobacco smoke compounds can directly affect the airway microbiota (e.g., enhanced biofilm formation by *Staphylococcus aureus*³⁶), and cigarettes themselves carry bacteria and fungi, including several human pathogens (Figure 1).³⁷ Furthermore, tobacco smoke upregulates the airway mucus production, impairs the mucociliary clearance, and induces low-grade inflammation within the lungs, collectively changing the micro-environmental conditions of the niche, which might affect the local microbiota.³⁸ These changes might eventually promote airway remodeling.^{39,40}

Many chronic lung diseases, including asthma and bronchiectasis, have been associated with an altered respiratory microbiota.^{34,41-43} The asthmatic bronchus with chronic inflammation and increased mucus production represents an ecological niche different from that of a healthy bronchus.³⁴ However, it is still unclear whether the bacterial dysbiosis is a cause or a consequence of the disease.⁴⁴ In this regard, the diverse compositions of the airway microbiota correlate

with the concentration of inflammatory cytokines in murine lungs, suggesting that changes in the microbiota can also modulate the host's inflammatory status.⁴⁵

2.2 | Other sources of indoor air pollution

Other agents, such as nitrogen dioxide (NO₂), carbon monoxide (CO), or some volatile organic compounds (VOC) (e.g., formaldehyde), are also main indoor pollutants.^{46,47}

Indoor NO₂ is mainly generated by gas-fueled cooking and heating appliances, and the levels in European households can be as high as 2500 µg/m³.⁴⁸ On the other hand, the *Towards Healthy Air in Dwellings in Europe* (THADE) project reported that the mean concentration of indoor NO₂ in Europe ranged from 10-15 µg/m³ in Scandinavia to 65 µg/m³ in Poland.⁴⁹ Numerous studies have reported positive associations between indoor NO₂ and the presence/aggravation of asthma symptoms in children.^{16,50} Some VOCs generated from sources like building materials or consumer products (cleaning products, cosmetics, air fresheners, etc) act as indoor air pollutants.⁵¹ A systematic review from 2015 reported only weak evidence relating exposure to this type of VOCs to the onset of asthma and AR, as well as to the aggravation of asthma/AR symptoms in both children and adults.⁵²

The use of solid fuel (e.g., coal) for cooking and heating does not only remain a source of indoor pollution in developing countries, but is still a common practice for residential heating in many Western households.⁵³ Exposure to the smoke generated by this biomass has




FIGURE 1 Tobacco smoke as driver of microbial dysbiosis in the airways: A, During homeostasis, there is a symbiosis between the airway microbiota and the stromal and immune cells of the respiratory epithelium; B, Tobacco smoke carries different microbes that can colonize the airways, and also promotes several changes in the resident microbiota such as the formation of biofilms by *Staphylococcus aureus*; C, Tobacco smoke upregulates mucus production, impairs the mucociliary clearance, and induces low-grade inflammation in the airway mucosa; C, The interaction between the effects of tobacco smoke on airway microbiota and respiratory epithelium further alters the ecological niche, promotes the outgrowth of certain species, and ultimately affects the microbial balance

been linked to several respiratory conditions in both adults and children,⁵⁴ but robust evidence is still lacking to support a causative role in the case of adult or pediatric asthma.

Indoor allergens from furry pets, molds, and house dust mites (HDM) also influence the quality of indoor air. Sensitization to furry animals is detected in up to 15% of the population⁵⁵ with a high degree of cross-reactivity among the different species. Moreover, HDM are the most common triggers of airway allergy, as up to 50% of asthmatics are sensitized to them.⁵⁶ Recent data suggest that in children with wheezing episodes, sensitization to HDM is associated with greater bronchial inflammation and reduced lung function.⁵⁷ Importantly, indoor allergens induce more severe phenotypes of airway allergy than outdoor seasonal allergens.⁵⁸⁻⁶⁰ Additionally, dampness is present in 10%-15% of households, which can lead to mold or cockroach colonization and subsequent allergic sensitization of the residents.^{61,62} Beyond allergic mechanisms, molds can promote inflammation of the upper and lower airways through several metabolites like glucans or mycotoxins.⁶³

In addition to residential environments, indoor air quality in nonresidential buildings (e.g., schools) plays an important role in respiratory health.⁶⁴⁻⁶⁶ The *Schools Indoor Pollution and Health: Observatory Network in Europe* (SINPHONIE) project, funded by the European Parliament, assessed indoor air exposure in schools in 23 countries.⁶⁷ The study found that PM_{2.5}, some VOCs (e.g., formaldehyde), radon, and allergens (especially molds) were commonly

present.⁶⁷ Moreover, the *Health Effects of School Environment* (HESE) project⁶⁸ reported that 78% and 66% of children attending schools in Norway, Sweden, Denmark, France, and Italy were exposed to PM₁₀ over 50 µg/m³ and to carbon dioxide (CO₂) over 1000 ppm, respectively. Another study from the HESE project investigating the burden of fungi showed that the number of viable molds in indoor air exceeded the maximum standard of 300 cfu/m³ in 33% of participating classrooms.⁶⁹ Very recently, these findings were confirmed in a study conducted in Southern Italy.⁷⁰ Importantly, this work observed that the concentration of elements from industrial emission was significantly higher in schools located in urban/industrial areas as compared to rural areas,⁷⁰ implying that the penetration of outdoor pollutants further deteriorates the quality of indoor air.

Table 1 summarizes the main effects of indoor pollution on AR and asthma.

3 | OUTDOOR AIR POLLUTION

3.1 | Epidemiological evidence

Various epidemiological studies have demonstrated that long-term exposure to outdoor air pollution (e.g., from traffic, industry) negatively affects respiratory health.⁷¹ A multicenter study in

TABLE 1 Consequences of indoor air pollution over allergic rhinitis and asthma

Indoor Air Pollution (effects amplified by the amount of time spent indoors and the isolation of buildings)	
Environmental factors	Health outcomes
Second-hand tobacco smoke	
During perinatal period	1. Higher prevalence of asthma, poorer asthma control, and more severe asthma exacerbations during childhood ¹⁹⁻²⁴ 2. Potential epigenetic changes with transgenerational repercussions ²⁵
During adulthood	1. Possibly higher asthma prevalence ²⁸⁻³⁰ 2. Possibly lower prevalence of allergic rhinitis ²⁹
Lifelong	Alteration of airway microbiota/bacterial dysbiosis ³⁴
NO ₂ from gas-fueled cooking and heating	Higher prevalence and exacerbation rate of asthma during childhood (16. 50)
Volatile organic compounds from building materials and consumer products	Possibly higher prevalence and exacerbation rate of allergic rhinitis and asthma in both children and adults ⁵²
Indoor allergens	
Lifelong	1. Increased sensitization rates to house dust mites, molds, and allergens from furry animals ^{55,56} 2. More severe phenotypes of allergic rhinitis and asthma, as compared to those induced by outdoor allergens ⁵⁸⁻⁶⁰
During childhood	Higher bronchial inflammation and reduced lung function in sensitized children with wheezing episodes ⁵⁷
Colonization of the households by molds and cockroaches due to dampness	1. Higher sensitization rate to dampness-related allergens ^{61,62} 2. Airway inflammation due to allergy-independent mechanisms (glucans, mycotoxins, etc) ⁶³

five European birth cohorts (conducted as part of the *European Study of Cohorts for Air Pollution Effects*, ESCAPE project) showed that exposure to PM and nitrogen oxides was associated with poor lung function in school-age children.⁷² An early analysis of the ESCAPE project did not find a statistically significant association between air pollution and the development of asthma up to the school age.⁷³ However, the re-analysis of these cohorts when the study individuals were 14-16 years old revealed associations between asthma incidence and NO₂ (OR 1.13, 95% CI: 1.02-1.25) and PM_{2.5} (OR 1.29, 95% CI: 1.00-1.66) exposure. The associations were particularly strong in the case of asthma diagnosed after the school age.⁷⁴ A recent meta-analysis of observational epidemiological studies published between 1999 and 2016 showed an association between traffic pollution and childhood asthma, with an OR ranging from 1.03 (95%CI: 1.01-1.05) to 1.08 (95%CI: 1.03-1.14) depending on the type of pollutant analyzed.⁷⁵ Other large studies have also reported similar links between outdoor pollution and childhood asthma.^{76,77}

The ESCAPE project also analyzed five European adult cohorts (overall ~7500 participants) and found that higher exposure to nitrogen monoxide (NO), NO₂, and PM₁₀ from traffic was associated with decreased lung function during adulthood.⁷⁸ A very recent study within the ESCAPE framework also found associations between decreased lung function and PM_{2.5} exposure in adults, with stronger effects observed for males.⁷⁹ Another work

within the ESCAPE project found suggestive (but nonsignificant) evidence that long-term exposure to NO₂, PM₁₀, and PM_{2.5} was associated with higher asthma incidence in adults.⁸⁰ Interestingly, a recent analysis of data from three large European cohort studies (with > 600 000 participants) showed that long-term PM₁₀ exposure was significantly associated with a 12.8% increase in lifetime asthma prevalence.⁸¹ Furthermore, several studies have confirmed the association between outdoor pollution and asthma exacerbations.⁸² Notably, a study from the *Improving Knowledge and Communication for Decision Making on Air Pollution and Health in Europe* (APHEKOM) network, which analyzed data from 10 European cities, showed that air pollution was accountable for up to 15% of all asthma exacerbations.⁸³

Regarding rhinitis, the urban dwelling has been related to a higher risk of AR as compared to the suburban dwelling.⁸⁴ Nevertheless, studies assessing the effect of air pollution on rhinitis onset have yielded inconsistent results in both children⁸⁵ and adults,⁸⁶ which contrasts with the large body of evidence relating air pollution to asthma.

Besides outdoor pollution from traffic, industry, energy production, heating, etc, emissions from livestock farming include specific pollutants such as organic dust, toxins from microorganisms, and gases like ammonia or methane.⁸⁷ These agents also influence the respiratory system either directly or through their role as precursors of other polluting particles. A large-scale population-based




FIGURE 2 Innate and adaptive immune effects of outdoor pollution on respiratory epithelium: A, PM₁₀ upregulates the expression of NLRP3 by airway epithelial cells (AEC). AECs in turn release CXCL1, which contributes to the recruitment of neutrophils from the bloodstream. AECs also release IL-6, which promotes mucus secretion; B, Black carbon induces the methylation and the expression of the IL-4 gene in different immune cells in the respiratory epithelium. Increased levels of IL-4 promote both mucosal Th2 cell priming and local IgE production by IgE + B cells. NLRP3: nucleotide-binding domain, leucine-rich repeat protein 3

study including ~2000 Dutch rural dwellers recently showed that increased levels of livestock-related air pollution were associated with decreased lung function, even in nonfarming individuals.⁸⁷

3.2 | Environmental-human interactions

The respiratory epithelium is composed of a pseudostratified layer of ciliated airway epithelial cells (AECs) intermingled with mucus-producing goblet cells.^{88,89} A recent study compared the response of primary murine and human AECs to either traffic-derived or ambient (collected in Sidney metropolitan area) PM_{2.5} and PM₁₀.⁹⁰ Noteworthy, ambient PM₁₀ induced a stronger secretion of IL-6 and CXCL1 by AECs, an effect attributed to the higher content of iron-rich particles from geological origin, as compared to traffic-derived PM₁₀. Importantly, PM₁₀-mediated secretion of cytokines is dependent on the *nucleotide-binding domain, leucine-rich repeat protein 3* (NLRP-3), a component of the inflammasome.⁹¹ The PM₁₀-mediated activation of the inflammasome induced profound innate immune effects in mouse models of allergic asthma, but was dispensable for PM₁₀-facilitated allergen sensitization.⁹¹ This finding indicates that PM₁₀ activates distinct inflammatory pathways, which might independently contribute to asthma pathogenesis.

Primary AECs from patients with severe asthma released more cytokines when exposed to PM or diesel exhaust (DE) compared to those from healthy subjects⁹² or patients with less severe asthma phenotypes.⁸⁹ This observation might explain how pollutant-induced epithelial insults^{93,94} can trigger asthma exacerbations.⁸³ Nevertheless, segmental allergen challenges in atopic subjects exposed to either DE or filtered air did not induce a different release of inflammatory mediators.^{89,95} This finding suggests that the adjuvant effect might rely on repetitive exposures. A recent mouse study⁹⁶ compared the effects of a two-hour exposure to DE or to the allegedly less toxic biodiesel (BD). Both exhaust products induced cardiovascular and pulmonary inflammation, while only BD generated an increase of neutrophils in bronchoalveolar lavage.

The environment acts on the genome inducing epigenetic changes, which function as important effectors of external insults. Epigenetic modification induces alterations in the DNA structure leading to changes in gene expression and inducing downstream disease.⁹⁷ Two recent studies analyzing pediatric populations from New York city showed a relationship between black carbon exposure and relevant epigenetic changes in immune genes.^{98,99} In a study analyzing samples from the oral mucosa, individuals with higher black carbon exposure had lower DNA methylation levels in the *IL-4* gene, possibly leading to higher expression.⁹⁹ This effect was even more significant in IgE-sensitized children⁹⁹ (Figure 2).

Importantly, most experimental studies apply pollutants in water solutions. This approach specifically selects for water-soluble components and modifies their chemical properties and interaction with AECs. This fact partly hampers the translation of the results from experimental settings to naturally occurring diseases. Moreover, in many studies, AECs are cultured submerged and in monolayers,

which does not reflect their natural physiology. Therefore, experimental designs using primary AECs grown at the air-liquid interface are required to investigate the real effect of PM or DE exposures on AEC responses in health and disease.

3.3 | Environment-environment interactions

Some air pollutants do not only have a direct effect on the respiratory system but also interact with plants and fungi to enhance the production and allergenicity of pollen, like ragweed or cypress,^{100,101} and of fungal spores.¹⁰² For example, ragweed in an urban area with high CO₂ concentrations grew faster and flowered earlier and more intensely, which led to the production of more pollen, as compared to ragweed grown in rural areas.¹⁰³ Recently, it has been shown that pollutants can also promote the release of allergens via direct cell damage.¹⁰⁴ Pollen grains and fungal spores contain several bioactive compounds, which may exert pro-inflammatory and pro-allergic effects.^{105,106} Recent data suggest that several pollen-associated lipid mediators (PALMs) activate Th2 cells to promote IgE synthesis in vitro.¹⁰⁷ Importantly, the pollen collected next to roads with heavy traffic released higher amounts of PALMs.¹⁰⁴ Similarly, a study carried out in Germany showed that birch trees exposed to higher concentrations of ozone produced more birch allergen (Bet v1) and PALMs per pollen grain than ozone-free trees.¹⁰⁸ Importantly, the skin prick test performed with an allergenic extract obtained from the highly exposed trees induced a significantly larger wheal diameter in birch pollen-AR patients as compared to the less exposed trees¹⁰⁸ (Figure 3).

Pollutants can also induce the oxidation or nitration of allergens, leading to changes in their conformation or stability. These chemical modifications enhance their immunogenicity and affect their interaction with receptors on immune cells.^{11,104} Several studies have shown that nitrated fungal spores¹⁰⁹ and nitrated pollen from birch, ragweed, and hornbeam^{108,110-112} have increased T cell-stimulatory and IgE-inducing capacity. Therefore, it is tempting to speculate that nitration by air pollutants plays a role in the IgE sensitization to allergens.¹⁰⁵

However, the clinical significance of these phenomena remains unclear.¹¹³ Epidemiological studies are largely limited by their inability to quantify individual exposure to air pollutants and allergens on a population scale. A French study including 36 397 AR patients found no effect of air pollution on the association between airborne pollen concentrations and rhinitis severity.¹¹⁴ Conversely, a study in eleven Canadian cities showed an interaction effect of air pollution on the risk of asthma-related hospitalizations and the burden of pollen grains and fungal spores.¹¹⁵ These findings have been recently confirmed in a study conducted in Belgium, which reported a synergistic effect of ozone, PM₁₀ and pollen exposure on the risk of asthma-related hospitalizations.¹¹⁶

Table 2 summarizes the main effects of outdoor pollution on AR and asthma.




FIGURE 3 Effect of outdoor pollution and climate change over allergenic plant species: Outdoor pollution increases the amount of pollen grains produced by plants (A), and the amount of both allergens (B) and PALMs (C) per pollen grain. Moreover, aeroallergens can become chemically modified by outdoor pollutants like nitrogen oxides (D). The global warming induces prolonged vegetation periods of allergenic plants (E), and extreme climate events like thunderstorms, which provoke the sudden release of massive amounts of allergens to the atmosphere (F). All these effects result on a higher availability of aeroallergens, and they increase the chances of interaction between the allergens and the stromal and immune cells of the airway mucosa. The interaction of native and nitrated allergens with airway epithelial cells can result on the release of pro-inflammatory mediators (G), whereas allergen interaction with dendritic cells can result on IgE sensitizations (H). The chances of sensitization are further increased by the higher availability of PALMs and nitrated allergens (G). Moreover, allergen interaction with sensitized mast cells can induce the release of inflammatory mediators (I), ultimately inducing the onset of asthma attacks in pollen-allergic patients (J). PALM: *pollen-associated lipid mediator*

4 | CLIMATE CHANGE

4.1 | Climate change and aeroallergens

Air pollution and climate change are closely interlinked. Nowadays, the vast majority of global energy is derived from fossil fuels whose burning generates huge amounts of CO₂, methane, black carbon, nitrogen oxides, and sulfate aerosols.¹¹⁷ Some of these pollutants (e.g., CO₂) are naturally occurring greenhouse gases, which persist for long periods in the atmosphere. Other agents (e.g., methane or black carbon) have shorter lifetimes but also contribute to climate change.¹¹⁸ Greenhouse gases help keep the earth warm by absorbing the sun's energy and by redirecting it back to the earth's surface.¹¹⁷ However, an overabundance of greenhouse gases traps an excessive amount of heat in the atmosphere and ultimately accounts for global warming.¹¹⁹

Global warming alters local vegetation patterns and speeds up the growth rate and phenology of plants, leading to increases in airborne pollen concentrations^{100,120} and changes in the geographical spread of plants.^{121,122} In this regard, climate change

was associated with increased duration of the ragweed pollen season in different studies conducted in North America and Europe.¹²³⁻¹²⁵ Changes in atmospheric humidity and precipitation also very likely affect the growth and distribution of fungi, yet this aspect remains uninvestigated.¹²⁶ The interactions of these changes with the photoperiod will modify the migration pattern of some plants and fungi.¹²⁷ Unlike air pollution,¹⁰⁸ global warming has not been related to date to enhanced allergenicity of plants.¹⁰⁰

The effects of climate change on allergenic plants and fungi¹²⁸⁻¹³⁰ are likely to continue in the future. A long-term prediction of these changes is challenging given the many variable factors, although computation efforts are currently ongoing.^{121,131,132} A process-based model of weed growth, plant competition, and population dynamics predicted that ragweed might spread to Northern European countries.¹²¹ Ragweed is a native species in North America, but is now rapidly invading several European areas.¹²¹ The colonization of geographical areas by new species will likely induce respiratory symptoms by both *de novo* sensitizations and cross-reactivity with pre-existing species.^{133,134}

TABLE 2 Consequences of outdoor air pollution over allergic rhinitis and asthma

Outdoor Air Pollution	
Environmental factors	Health outcomes
Pollution from traffic and industry (PM ₁₀ , PM _{2.5} , NO, NO ₂ , etc)	
During childhood	Higher asthma prevalence after the school age ⁷⁴⁻⁷⁷
During adulthood	Possibly higher asthma prevalence ^{80,81}
Lifelong	1. Poorer lung function ^{72,78,79} 2. Higher rate of asthma exacerbations (82. 83) 3. Conflicting results on AR onset ^{85,86}
Livestock farming (organic dust, toxins from microorganisms, gases like ammonia and methane, etc)	Decreased lung function ⁸⁷
Black carbon	Possibly epigenetic changes leading to increased type two inflammation in children ⁹⁹
Interaction between air pollutants (PM ₁₀ , nitrogen oxides) and allergens (pollen, fungal spores, etc)	
Production of more pollen, more allergens per pollen grain, and more PALMs per pollen grain ^{100-104,108}	1. Potentially, facilitation of IgE sensitization against aeroallergens ^{104,107} 2. Higher rate of asthma-related hospitalizations ^{115,116}
Release of allergens via direct cell damage ¹⁰⁴	
Nitration of allergens ¹⁰⁹⁻¹¹²	

Abbreviation: PALM, pollen-associated lipid mediator.

4.2 | Climate change as an inducer of respiratory and allergic diseases

There is no doubt that climate change causes or exacerbates respiratory diseases.¹³⁵⁻¹³⁹ The most important effects of climate change on respiratory health are described below (also summarized in Table 3):

1. The higher temperatures and increased frequency of heatwaves amplify the exacerbation rate, morbidity, and mortality of respiratory diseases.¹⁴⁰⁻¹⁴³ The extent of this association usually parallels the pollution levels of local air.¹⁴⁴
2. The seasonality and severity of AR and asthma are affected by the growth patterns of allergenic species,¹⁴⁵⁻¹⁴⁸ which can act synergistically with air pollutants.¹⁴⁹ Global warming might also alter the species dominating distinct ecological niches.¹⁵⁰
3. Climate change is expected to alter the pattern of respiratory tract infections.^{151,152}
4. Intensive rain and flooding induce dampness and mold proliferation in affected households,^{61,153} thus influencing the quality of indoor air.

5. Extreme climate events are the cause of specific phenomena like thunderstorm-related asthma episodes.^{153,154} During these episodes, a large number of patients experience asthmatic symptoms during the initial 20-30 minutes of a large-scale thunderstorm, providing that it occurs during the allergen season and induces a cold outflow.¹⁵⁵ This phenomenon arises from a sudden release of massive amounts of aeroallergens,¹⁵⁴ and a causative role for pollen allergy is suspected.¹⁵⁶ Numerous case studies of thunderstorm-related asthma have been documented,⁶¹ the largest of which took place in Melbourne (Australia) on November 21, 2016 (~4000 patients presented at hospitals with respiratory symptoms).^{156,157}

Besides the direct effects of global warming on the airways, the altered levels of aeroallergens account from many of the effects of climate change on respiratory health. As climate change will also influence the amount and type of pollutants in the air, which themselves interact with aeroallergens,¹³⁸ the individual and/or combined effects of these environmental parameters on respiratory health are very difficult to predict.

5 | INTERVENTIONS TO MODIFY AIR POLLUTION AND CLIMATE CHANGE

5.1 | Policy changes

Policy changes are the most effective measures to decrease pollution.^{158,159} While actions of individual citizens can mitigate air pollution only to a small extent,¹⁶⁰ larger lifestyle changes at the population level mainly result from policy interventions. For example, many countries have implemented smoke-free legislation to protect the population, particularly children in public places. A recent meta-analysis of 35 pediatric studies showed that enforcement of smoke-free policies was significantly associated with a 9.8% (95%CI: 3%-16%) and 18.5% (95%CI: 4.2%-32.8%) reduction of hospital admissions due to asthma attacks and lower respiratory tract infections, respectively.¹³ These associations tended to be stronger in regions with more comprehensive smoke-free laws, indicating that stringent smoke-free policies are necessary to gain maximum health benefits.

The replacement of fossil fuels by renewable energy sources and commitment to a complete phaseout of coal power by the industry represent necessary milestones in the roadmap for a more environmentally friendly economy.^{161,162} Over the past decade, the implementation of the European Union (EU) environmental policy framework contributed substantially to decreasing the emissions of many air pollutants and improving air quality across Europe.¹⁶³ The EU recently released an updated version of the *environmental performance standards for large combustion plants*, which set stricter emission ranges for NO, NO₂, sulfur dioxide, PM, and mercury from power plants.¹⁶⁴ A recent *National Emission Ceiling Directive* was also released as a measure to reduce emissions from different sectors.¹⁶⁵

Environmental changes	Health outcomes
More frequent extreme climate events	
Heatwaves, wildfires, higher temperatures, etc	Amplification of exacerbation rate, morbidity, and mortality of respiratory diseases. ¹⁴⁰⁻¹⁴³
Intensive rain and flooding	Dampness in affected households with subsequent proliferation of molds and cockroaches. ^{61,153} See the consequences of the deterioration of indoor air quality in Table 1.
Thunderstorms	Increase in asthma exacerbations and hospitalizations following thunderstorm-related asthma episodes ¹⁵³⁻¹⁵⁷
More intense and more prolonged pollen seasons. ^{100,120,123-125} Possibly similar changes for other allergens (e.g., fungi).	Increase in the severity and alteration of the seasonality of symptoms of allergic rhinitis and asthma ¹⁴⁵⁻¹⁴⁸
Alteration of the local vegetation patterns with changes in the geographical spread ¹²² and migration of plants ¹²⁷ Colonization of geographical areas by new species ¹²¹ with alteration of the species dominating distinct ecological niches. ¹⁵⁰ Possibly, similar changes for fungi. ¹²⁶	Increased prevalence and severity of allergic rhinitis and asthma due to both de novo sensitizations and cross-reactivity with pre-existing species ^{133,134}
Possibly changes in the growth pattern and distribution of pathogenic microorganisms. ¹⁵¹	Possibly changes in the pattern of respiratory tract infections

TABLE 3 Consequences of climate change and global warming over allergic rhinitis and asthma

Nevertheless, a recent report from the *European Economic Area* showed that a large proportion of European citizens and ecosystems are still exposed to concentrations of air pollutants exceeding the legal limit values of the EU and the guideline values of the WHO (Table 4).² Renewable energy sources currently account for 24% of total electricity generated,¹⁶⁶ illustrating the long way to go before fossil fuels can be replaced. Government investment in clean energy should come together with policies incenting suppliers into a timely transition out of existing fossil-based infrastructure.¹⁶⁶

At a local level, greenhouse-gas emissions can be reduced by shifting from private motorized transport to more sustainable modalities, such as public transport, cycling, and walking. There is evidence that having good cycling infrastructure integrated with public transport, training of both cyclists and motorists, and making driving costly can promote cycling.¹⁶⁶ Local authorities could also incentivize the population to shift to sustainable electric vehicles by introducing ownership tax exemptions and additional advantages such as waivers on fees (e.g., plug-in charging station or parking spots). Studies evaluating the effect of nearby green areas on respiratory health have yielded inconsistent results,¹⁶⁷ possibly due to the complex interactions between global warming, vegetation, and air pollution. Recent studies using the methodology recommended by the *Coordination of Information on the Environment* (CORINE) program have yielded conflicting results, indicating either a beneficial impact of greenness on general health,¹⁶⁸ or an increased risk of wheezing, asthma, and AR in children exposed to green spaces, especially coniferous forests.¹⁶⁹ Until future studies shed more light on this issue, it seems reasonable that city development plans include green spaces with diverse and nonallergenic species.^{170,171}

The *European Academy of Allergy and Clinical Immunology* (EAACI)/*European Federation for Allergy and Airways Diseases Patients' Associations* (EFA) Interest Group on Allergy and Asthma at the European Parliament recently launched a call to increase the awareness about asthma and promote the legal changes required to decrease the burden of air pollution.^{172,173}

5.2 | Lifestyle adjustments

Various lifestyle adjustments can mitigate air pollution and climate change and indirectly decrease the onset and progression of respiratory diseases. The avoidance of individual motorized transportation constitutes a simple and basic approach. Exercising outdoors is also recommended, as its benefits are expected to exceed the negative impacts of exposure to outdoor allergens and pollutants, at least in most European cities.^{174,175} However, limiting the time spent outdoors during the pollen season (for pollen-allergic patients)¹⁷⁶ and during high traffic hours or warm days is a reasonable approach. Air quality alerts, pollen calendars, and allergy diaries, among other mobile health tools, can help plan outdoor activities, and control and monitor symptoms.^{177,178} Besides mobility, livestock to provide meat for human consumption is among the main causes of CO₂ production. Therefore, reduction of meat consumption, together with limiting trips made by air traffic, and increasing the use of recyclable materials, are also meaningful measures to reduce CO₂ emissions.

Adequate and regular ventilation of the living spaces and the filtration of indoor air can prevent mold growth and reduce indoor NO₂ concentrations. Moreover, preventing dampness

TABLE 4 Percentage of the urban population in the EU-28 exposed to air pollutant concentrations above the EU limit or target values and above the WHO air quality guidelines (AQGs)

Pollutant	EU reference value	Exposure estimate (%)	WHO AQG	Exposure estimate (%)
PM _{2.5}	Year (25)	7-8	Year (10)	82-85
PM ₁₀	Day (50)	16-20	Year (20)	50-62
O ₂	8-h (120)	7-30	8-h (100)	95-98
NO ₂	Year (40)	7-9	Year (40)	7-9
BaP	Year (1)	20-25	Year (0.12) RL	85-91
SO ₂	Day (125)	<1	Day (20)	20-38
Key	<5%	5%-50%	50%-75%	>75%

Note: Minimum and maximum values observed between 2013 and 2015. The comparison is made for the most stringent EU limit or target values set for the protection of human health. As the WHO has not set AQGs for benzo(a)pyrene (BaP), the reference level in the table was estimated assuming WHO unit risk for lung cancer for PAH mixtures and an acceptable risk of additional lifetime cancer risk of approximately 1 in 100 000. Estimated reference levels are given between parentheses and in $\mu\text{g}/\text{m}^3$, except for BaP, which is given in ng/m^3 . This table was modified and reproduced with permission from the European Environmental Agency Report No 13/2017. Air quality in Europe—2017 report. Available from: <https://www.eea.europa.eu/publications/air-quality-in-europe-2017> (assessed: July 2018).

and limiting the number of carpets can decrease the burden of HDMs and molds.¹⁷⁹ Nevertheless, in areas with a high burden of HDMs, these strategies are often insufficient, as emphasized by international guidelines and consensus documents on AR and asthma.¹⁸⁰⁻¹⁸² Moreover, individuals with an atopic predisposition should carefully consider the choice of keeping a pet.^{179,183} In this regard, a large epidemiological survey identified exposure to cat during the first year of life as an independent risk factor for AR and asthma presence during school age.¹⁸⁴

6 | CONCLUSION

Given the explosive global rise in urbanization, industrial production, aviation, road traffic, etc, the preservation of good air quality will become increasingly challenging. This narrative review summarizes the current literature about the potential effects of air pollution and climate change on AR and asthma. Although a formal meta-analysis was outside the scope of this review, and there were several difficulties for direct comparisons of the studies due to methodological differences, several conclusions could be made.

1. The evidence relating passive smoking and exposure to traffic-related pollution (including NO₂ and PM_{2.5}) to childhood asthma is currently robust, whereas the link with exposure to smoke from biomass and deleterious VOCs seems weaker.
2. Although the relationship between air pollution and adult-onset asthma has been uncertain for years, recent data suggest that passive smoking and traffic pollutants might be related to asthma development in adults.
3. The relationship between air pollution and AR onset seems less conclusive as compared to asthma in both children and adults.

4. The mechanisms for how pollutants induce respiratory disease are varied. Recent evidence indicates that epigenetic changes in the respiratory epithelium and the alteration of airway microbiota might account for some of the effects of PM_{2.5} and tobacco smoke, respectively.
5. Exposure to indoor and outdoor allergens is a well-established risk factor for the development of AR and asthma in both adults and children, with indoor allergens inducing more severe phenotypes of airway allergy.
6. The capacity of outdoor pollutants to increase the allergenicity and immunogenicity of aeroallergens has been shown in vitro, but the clinical implications of these phenomena require further analysis.
7. Unlike pollutants, climate change affects pollen grains and fungal spores by increasing their availability rather than altering their chemical structure. Some of the deleterious effects of climate change on respiratory health are likely to arise from this increased availability.

As a key message, we can conclude that the detrimental effects of air pollution and climate change on human health are greatly preventable through timely implementation of adequate legislations. Governments need to adopt effective and evidence-based regulations, as political interventions are the only way to achieve large improvements at the population level. All these efforts are crucial steps in the pathway to clean air, and ultimately, to the prevention and reduction of AR, asthma, and other chronic respiratory conditions.

ACKNOWLEDGMENT

IEG receives funding from Instituto de Salud Carlos III, Spanish Ministry of Science and Innovation through the Rio Hortega and

RETICS schemes (CM17/00140 and RD16/0006/0001). AGM and JV are supported by the National Institute of Health Research Manchester Biomedical Research Centre (NIHR Manchester BRC). Y.C is supported by a Medical Research Council Early Career Research Fellowship awarded through the MRC-PHE Centre for Environment and Health (grant number MR/M501669/1).










CONFLICTS OF INTEREST

Apart from academic affiliations, ZD acts as Executive and Scientific Medical Director at a phase I/II pharmacological unit (QPS-NL), which performs clinical studies for pharmaceutical companies. In the past 3 years, ZD received honoraria, consultancy and speaker fees from Astrazeneca, ALK, Aquilon, Boehringer Ingelheim, CSL, HAL Allergy, MSD, Sanofi-Genzyme. The remaining authors have no conflicts of interest to declare.

AUTHOR CONTRIBUTIONS

IEG and AM coordinated the work and designed the structure of the review. SB, SJHV, EF, PC, YSC, and PVT reviewed the literature, wrote the manuscript, and prepared the figures and tables. IEG, AM, ZD, JV, CG, and BH critically reviewed and finalized the manuscript.

ORCID

Ibon Eguluz-Gracia  <https://orcid.org/0000-0002-3774-931X>
 Alexander G. Mathioudakis  <https://orcid.org/0000-0002-4675-9616>
 Sabine Bartel  <https://orcid.org/0000-0002-9163-795X>
 Susanne J. H. Vijverberg  <https://orcid.org/0000-0002-4579-4081>
 Elaine Fuertes  <https://orcid.org/0000-0003-0205-9025>
 Peter Valentin Tomazic  <https://orcid.org/0000-0001-6445-4800>
 Zuzana Diamant  <https://orcid.org/0000-0003-0133-0100>
 Jørgen Vestbo  <https://orcid.org/0000-0001-6355-6362>
 Carmen Galan  <https://orcid.org/0000-0002-6849-1219>

REFERENCES

- World Health Organisation Report. 9 out of 10 people worldwide breathe polluted air, but more countries are taking action. <http://www.who.int/news-room/detail/02-05-2018-9-out-of-10-people-worldwide-breathe-polluted-air-but-more-countries-are-taking-action>. Accessed July 1, 2018.
- Air quality in Europe – 2018 report. <https://www.eea.europa.eu/publications/air-quality-in-europe-2018>. Accessed July 1, 2018.
- Gakidou E, Afshin A, Abajobir AA. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;390(10100):1345-1422.
- Greenland S. Concepts and pitfalls in measuring and interpreting attributable fractions, prevented fractions, and causation probabilities. *Ann Epidemiol*. 2015;25(3):155-161.
- Del Giacco SR, Bakirtas A, Bel E, et al. Allergy in severe asthma. *Allergy*. 2017;72(2):207-220.
- Rondon C, Bogas G, Barrionuevo E, Blanca M, Torres MJ, Campo P. Nonallergic rhinitis and lower airway disease. *Allergy*. 2017;72(1):24-34.
- Colas C, Brosa M, Anton E, et al. Estimate of the total costs of allergic rhinitis in specialized care based on real-world data: the FERIN Study. *Allergy*. 2017;72(6):959-966.
- Belhassen M, Demoly P, Bloch-Morot E, et al. Costs of perennial allergic rhinitis and allergic asthma increase with severity and poor disease control. *Allergy*. 2017;72(6):948-958.
- Bousquet J, Khaltaev N, Cruz AA, et al. Allergic rhinitis and its impact on asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy*. 2008;63(Suppl 86):8-160.
- Agache I, Miller R, Gern JE, et al. Emerging concepts and challenges in implementing the exposome paradigm in allergic diseases and asthma: a Practall document. *Allergy*. 2019;74(3):449-463.
- Cecchi L, D'Amato G, Annesi-Maesano I. External exposome and allergic respiratory and skin diseases. *J Allergy Clin Immunol*. 2018;141(3):846-857.
- Abramson MJ, Guo Y. Indoor endotoxin exposure and ambient air pollutants interact on asthma outcomes. *Am J Respir Crit Care Med*. 2019;200(6):652-654.
- Faber T, Kumar A, Mackenbach JP, et al. Effect of tobacco control policies on perinatal and child health: a systematic review and meta-analysis. *Lancet Public Health*. 2017;2(9):e420-e437.
- Molloy J, Koplin JJ, Allen KJ, et al. Vitamin D insufficiency in the first 6 months of infancy and challenge-proven IgE-mediated food allergy at 1 year of age: a case-cohort study. *Allergy*. 2017;72(8):1222-1231.
- Yepes-Nunez JJ, Brozek JL, Fiocchi A, et al. Vitamin D supplementation in primary allergy prevention: Systematic review of randomized and non-randomized studies. *Allergy*. 2018;73(1):37-49.
- Breyse PN, Diette GB, Matsui EC, Butz AM, Hansel NN, McCormack MC. Indoor air pollution and asthma in children. *Proc Am Thorac Soc*. 2010;7(2):102-106.
- Hulin M, Simoni M, Viegi G, Annesi-Maesano I. Respiratory health and indoor air pollutants based on quantitative exposure assessments. *Eur Respir J*. 2012;40(4):1033-1045.
- Drago G, Perrino C, Canepari S, et al. Relationship between domestic smoking and metals and rare earth elements concentration in indoor PM_{2.5}. *Environ Res*. 2018;165:71-80.
- Vanker A, Gie RP, Zar HJ. The association between environmental tobacco smoke exposure and childhood respiratory disease: a review. *Expert Rev Respir Med*. 2017;11(8):661-673.
- Vardavas CI, Hohmann C, Patelarou E, et al. The independent role of prenatal and postnatal exposure to active and passive smoking on the development of early wheeze in children. *Eur Respir J*. 2016;48(1):115-124.
- Schwarze J, Openshaw P, Jha A, et al. Influenza burden, prevention, and treatment in asthma-A scoping review by the EAACI Influenza in asthma task force. *Allergy*. 2018;73(6):1151-1181.
- Feldman LY, Thacher JD, vanHage M, et al. Early-life secondhand smoke exposure and food hypersensitivity through adolescence. *Allergy*. 2018;73(7):1558-1561.
- Burke H, Leonardi-Bee J, Hashim A, et al. Prenatal and passive smoke exposure and incidence of asthma and wheeze: systematic review and meta-analysis. *Pediatrics*. 2012;129(4):735-744.
- Thacher JD, Gehring U, Gruzieva O, et al. Maternal smoking during pregnancy and early childhood and development of asthma and rhinoconjunctivitis - a MeDALL project. *Environ Health Perspect*. 2018;126(4):047005.
- Melen E, Barouki R, Barry M, et al. Promoting respiratory public health through epigenetics research: an ERS Environment Health Committee workshop report. *Eur Respir J*. 2018;51(4):1702410.
- Hwang SH, Hwang JH, Moon JS, Lee DH. Environmental tobacco smoke and children's health. *Korean J Pediatr*. 2012;55(2):35-41.
- Accordini S, Janson C, Svanes C, Jarvis D. The role of smoking in allergy and asthma: lessons from the ECRHS. *Curr Allergy Asthma Rep*. 2012;12(3):185-191.

28. Coogan PF, Castro-Webb N, Yu J, O'Connor GT, Palmer JR, Rosenberg L. Active and passive smoking and the incidence of asthma in the Black Women's Health Study. *Am J Respir Crit Care Med*. 2015;191(2):168-176.
29. Skaaby T, Taylor AE, Jacobsen RK, et al. Investigating the causal effect of smoking on hay fever and asthma: a Mendelian randomization meta-analysis in the CARTA consortium. *Sci Rep*. 2017;7(1):2224.
30. Flexeder C, Zock JP, Jarvis D, et al. Second-hand smoke exposure in adulthood and lower respiratory health during 20 year follow up in the European Community Respiratory Health Survey. *Respir Res*. 2019;20(1):019-0996.
31. Simonyte Sjodin K, Hammarstrom ML, Ryden P, et al. Temporal and long-term gut microbiota variation in allergic disease: a prospective study from infancy to school age. *Allergy*. 2019;74(1):176-185.
32. Sbihi H, Boutin RC, Cutler C, Suen M, Finlay BB, Turvey SE. Thinking bigger: How early-life environmental exposures shape the gut microbiome and influence the development of asthma and allergic disease. *Allergy*. 2019;9(10):13812.
33. Birzele LT, Depner M, Ege MJ, et al. Environmental and mucosal microbiota and their role in childhood asthma. *Allergy*. 2017;72(1):109-119.
34. Marsland BJ, Gollwitzer ES. Host-microorganism interactions in lung diseases. *Nat Rev Immunol*. 2014;14(12):827-835.
35. Morris A, Beck JM, Schloss PD, et al. Comparison of the respiratory microbiome in healthy nonsmokers and smokers. *Am J Respir Crit Care Med*. 2013;187(10):1067-1075.
36. Kulkarni R, Antala S, Wang A, et al. Cigarette smoke increases *Staphylococcus aureus* biofilm formation via oxidative stress. *Infect Immun*. 2012;80(11):3804-3811.
37. Sapkota AR, Berger S, Vogel TM. Human pathogens abundant in the bacterial metagenome of cigarettes. *Environ Health Perspect*. 2010;118(3):351-356.
38. Jaspers I. Cigarette smoke effects on innate immune mechanisms in the nasal mucosa. Potential effects on the microbiome. *Ann Am Thorac Soc*. 2014;11(Suppl 1):S38-S42.
39. Samitas K, Carter A, Kariyawasam HH, Xanthou G. Upper and lower airway remodelling mechanisms in asthma, allergic rhinitis and chronic rhinosinusitis: the one airway concept revisited. *Allergy*. 2018;73(5):993-1002.
40. Mehta AK, Doherty T, Broide D, Croft M. Tumor necrosis factor family member LIGHT acts with IL-1beta and TGF-beta to promote airway remodeling during rhinovirus infection. *Allergy*. 2018;73(7):1415-1424.
41. Lee JJ, Kim SH, Lee MJ, et al. Different upper airway microbiome and their functional genes associated with asthma in young adults and elderly individuals. *Allergy*. 2019;74(4):709-719.
42. Lunjani N, Satitsuksano P, Lukasik Z, Sokolowska M, Eiwegger T, O'Mahony L. Recent developments and highlights in mechanisms of allergic diseases: microbiome. *Allergy*. 2018;73(12):2314-2327.
43. Simpson JL, Daly J, Baines KJ, et al. Airway dysbiosis: *Haemophilus influenzae* and *Tropheryma* in poorly controlled asthma. *Eur Respir J*. 2016;47(3):792-800.
44. Kim BS, Lee E, Lee MJ, et al. Different functional genes of upper airway microbiome associated with natural course of childhood asthma. *Allergy*. 2018;73(3):644-652.
45. Dickson RP, Erb-Downward JR, Falkowski NR, Hunter EM, Ashley SL, Huffnagle GB. The lung microbiota of healthy mice are highly variable, cluster by environment, and reflect variation in baseline lung innate immunity. *Am J Respir Crit Care Med*. 2018;198(4):497-508.
46. Azizi BH, Zulkifli HI, Kasim S. Indoor air pollution and asthma in hospitalized children in a tropical environment. *J Asthma*. 1995;32(6):413-418.
47. Logue JM, Klepeis NE, Lobscheid AB, Singer BC. Pollutant exposures from natural gas cooking burners: a simulation-based assessment for Southern California. *Environ Health Perspect*. 2014;122(1):43-50.
48. Jarvis DJAG, Heroux ME, Rapp R, Kelly FK. Nitrogen dioxide. In: WHO Guidelines for Indoor Air Quality: Selected Pollutants. Geneva: World Health Organization; 2010. 5. <https://www.ncbi.nlm.nih.gov/books/NBK138707/>. Accessed July 1, 2018.
49. Franchi M, et al. Towards healthy air in dwellings in Europe. Brussels: European Federation of Allergy and Airways Diseases Patients Associations. 2004.
50. Gillespie-Bennett J, Piersie N, Wickens K, Crane J, Howden-Chapman P. The respiratory health effects of nitrogen dioxide in children with asthma. *Eur Respir J*. 2011;38(2):303-309.
51. United States Environmental Protection Agency. An Introduction to Indoor Air Quality (IAQ). Volatile Organic Compounds (VOCs). www.epa.gov/iaq/voc.html. Accessed July 9, 2012.
52. Nurmatov UB, Tagiyeva N, Semple S, Devereux G, Sheikh A. Volatile organic compounds and risk of asthma and allergy: a systematic review. *Eur Respir Rev*. 2015;24(135):92-101.
53. Residential heating with wood and coal: health impacts and policy options in Europe and North America. http://www.euro.who.int/_data/assets/pdf_file/0009/271836/ResidentialHeatingWoodCoalHealthImpacts.pdf
54. Kurmi OP, Lam KB, Ayres JG. Indoor air pollution and the lung in low- and medium-income countries. *Eur Respir J*. 2012;40(1):239-254.
55. Brunekreef B, Von Mutius E, Wong G, Odhiambo J, Garcia-Marcos L, Foliaki S. Exposure to cats and dogs, and symptoms of asthma, rhinoconjunctivitis, and eczema. *Epidemiology*. 2012;23(5):742-750.
56. de Vries MP, van den Bemt L, van der Mooren FM, Muris JW, van Schayck CP. The prevalence of house dust mite (HDM) allergy and the use of HDM-impermeable bed covers in a primary care population of patients with persistent asthma in the Netherlands. *Prim Care Respir J*. 2005;14(4):210-214.
57. Ruggieri S, Drago G, Longo V, et al. Sensitization to dust mite defines different phenotypes of asthma: a multicenter study. *Pediatr Allergy Immunol*. 2017;28(7):675-682.
58. Platts-Mills TA. The allergy epidemics: 1870-2010. *J Allergy Clin Immunol*. 2015;136(1):3-13.
59. Cazzoletti L, Marcon A, Corsico A, et al. Asthma severity according to Global Initiative for Asthma and its determinants: an international study. *Int Arch Allergy Immunol*. 2010;151(1):70-79.
60. Calderon MA, Linneberg A, Kleine-Tebbe J, et al. Respiratory allergy caused by house dust mites: What do we really know? *J Allergy Clin Immunol*. 2015;136(1):38-48.
61. Katelaris CH, Beggs PJ. Climate change: allergens and allergic diseases. *Intern Med J*. 2018;48(2):129-134.
62. Thacher JD, Gruzieva O, Pershagen G, et al. Mold and dampness exposure and allergic outcomes from birth to adolescence: data from the BAMSE cohort. *Allergy*. 2017;72(6):967-974.
63. Flamant-Hulin M, Annesi-Maesano I, Caillaud D. Relationships between molds and asthma suggesting non-allergic mechanisms. A rural-urban comparison. *Pediatr Allergy Immunol*. 2013;24(4):345-351.
64. Takaoka M, Suzuki K, Norback D. Current asthma, respiratory symptoms and airway infections among students in relation to the school and home environment in Japan. *J Asthma*. 2017;54(6):652-661.
65. Gaffin JM, Hauptman M, Petty CR, et al. Nitrogen dioxide exposure in school classrooms of inner-city children with asthma. *J Allergy Clin Immunol*. 2018;141(6):2249-2255.e2.
66. Sander I, Lotz A, Neumann HD, et al. Indoor allergen levels in settled airborne dust are higher in day-care centers than at home. *Allergy*. 2018;73(6):1263-1275.
67. Csobod EA-MI, Carrer P, et al. SINFONIE - Schools Indoor Pollution and Health Observatory Network in Europe - Final

- Report. Publications Office of the European Union; 2014. <http://publications.jrc.ec.europa.eu/repository/handle/JRC91160>. Accessed June 8, 2018.
68. Simoni M, Annesi-Maesano I, Sigsgaard T, et al. School air quality related to dry cough, rhinitis and nasal patency in children. *Eur Respir J*. 2010;35(4):742-749.
 69. Simoni M, Cai GH, Norback D, et al. Total viable molds and fungal DNA in classrooms and association with respiratory health and pulmonary function of European schoolchildren. *Pediatr Allergy Immunol*. 2011;22(8):843-852.
 70. Ruggieri S, Longo V, Perrino C, et al. Indoor air quality in schools of a highly polluted south Mediterranean area. *Indoor Air*. 2019;29(2):276-290.
 71. Thurston GD, Kipen H, Annesi-Maesano I, et al. A joint ERS/ATS policy statement: what constitutes an adverse health effect of air pollution? An analytical framework. *Eur Respir J*. 2017;49(1):1600419.
 72. Gehring U, Gruzieva O, Agius RM, et al. Air pollution exposure and lung function in children: the ESCAPE project. *Environ Health Perspect*. 2013;121(11-12):1357-1364.
 73. Molter A, Simpson A, Berdel D, et al. A multicentre study of air pollution exposure and childhood asthma prevalence: the ESCAPE project. *Eur Respir J*. 2015;45(3):610-624.
 74. Gehring U, Wijga AH, Hoek G, et al. Exposure to air pollution and development of asthma and rhinoconjunctivitis throughout childhood and adolescence: a population-based birth cohort study. *Lancet Respir Med*. 2015;3(12):933-942.
 75. Khreis H, Kelly C, Tate J, Parslow R, Lucas K, Nieuwenhuijsen M. Exposure to traffic-related air pollution and risk of development of childhood asthma: a systematic review and meta-analysis. *Environ Int*. 2017;100:1-31.
 76. Hsu HH, Chiu YH, Coull BA, et al. Prenatal particulate air pollution and asthma onset in urban children. Identifying sensitive windows and sex differences. *Am J Respir Crit Care Med*. 2015;192(9):1052-1059.
 77. Sbihi H, Tamburic L, Koehoorn M, Brauer M. Perinatal air pollution exposure and development of asthma from birth to age 10 years. *Eur Respir J*. 2016;47(4):1062-1071.
 78. Adam M, Schikowski T, Carsin AE, et al. Adult lung function and long-term air pollution exposure. ESCAPE: a multicentre cohort study and meta-analysis. *Eur Respir J*. 2015;45(1):38-50.
 79. Doiron D, de Hoogh K, Probst-Hensch N, et al. Air pollution, lung function and COPD: results from the population-based UK Biobank study. *Eur Respir J*. 2019;54(1):1802140.
 80. Jacquemin B, Siroux V, Sanchez M, et al. Ambient air pollution and adult asthma incidence in six European cohorts (ESCAPE). *Environ Health Perspect*. 2015;123(6):613-621.
 81. Cai Y, Zijlema WL, Doiron D, et al. Ambient air pollution, traffic noise and adult asthma prevalence: a BioSHaRE approach. *Eur Respir J*. 2017;49(1):1502127.
 82. Tanaka H, Nakatani E, Fukutomi Y, et al. Identification of patterns of factors preceding severe or life-threatening asthma exacerbations in a nationwide study. *Allergy*. 2018;73(5):1110-1118.
 83. Perez L, Declercq C, Iniguez C, et al. Chronic burden of near-roadway traffic pollution in 10 European cities (APHEKOM network). *Eur Respir J*. 2013;42(3):594-605.
 84. Maio S, Baldacci S, Carrozzi L, et al. Respiratory symptoms/diseases prevalence is still increasing: a 25-yr population study. *Respir Med*. 2016;110:58-65.
 85. Deng Q, Lu C, Yu Y, Li Y, Sundell J, Norback D. Early life exposure to traffic-related air pollution and allergic rhinitis in preschool children. *Respir Med*. 2016;121:67-73.
 86. Burte E, Leynaert B, Bono R, et al. Association between air pollution and rhinitis incidence in two European cohorts. *Environ Int*. 2018;115:257-266.
 87. Borlee F, Yzermans CJ, Aalders B, et al. Air pollution from livestock farms is associated with airway obstruction in neighboring residents. *Am J Respir Crit Care Med*. 2017;196(9):1152-1161.
 88. Whitsett JA, Alenghat T. Respiratory epithelial cells orchestrate pulmonary innate immunity. *Nat Immunol*. 2015;16(1):27-35.
 89. Weng CM, Wang CH, Lee MJ, et al. Aryl hydrocarbon receptor activation by diesel exhaust particles mediates epithelium-derived cytokines expression in severe allergic asthma. *Allergy*. 2018;73(11):2192-2204.
 90. Kumar RK, Shadie AM, Bucknall MP, et al. Differential injurious effects of ambient and traffic-derived particulate matter on airway epithelial cells. *Respirology*. 2015;20(1):73-79.
 91. Hirota JA, Gold MJ, Hiebert PR, et al. The nucleotide-binding domain, leucine-rich repeat protein 3 inflammasome/IL-1 receptor 1 axis mediates innate, but not adaptive, immune responses after exposure to particulate matter under 10 μm . *Am J Respir Cell Mol Biol*. 2015;52(1):96-105.
 92. Hackett TL, Singhera GK, Shaheen F, et al. Intrinsic phenotypic differences of asthmatic epithelium and its inflammatory responses to respiratory syncytial virus and air pollution. *Am J Respir Cell Mol Biol*. 2011;45(5):1090-1100.
 93. Jain V, Raina S, Gheware AP, et al. Reduction in polyamine catabolism leads to spermine-mediated airway epithelial injury and induces asthma features. *Allergy*. 2018;73(10):2033-2045.
 94. Mitamura Y, Nunomura S, Nanri Y, et al. The IL-13/periostin/IL-24 pathway causes epidermal barrier dysfunction in allergic skin inflammation. *Allergy*. 2018;73(9):1881-1891.
 95. Rider CF, Yamamoto M, Gunther OP, et al. Controlled diesel exhaust and allergen coexposure modulates microRNA and gene expression in humans: Effects on inflammatory lung markers. *J Allergy Clin Immunol*. 2016;138(6):1690-1700.
 96. de Brito JM, Mauad T, Cavalheiro GF, et al. Acute exposure to diesel and sewage biodiesel exhaust causes pulmonary and systemic inflammation in mice. *Sci Total Environ*. 2018;629:1223-1233.
 97. Portela A, Esteller M. Epigenetic modifications and human disease. *Nat Biotechnol*. 2010;28(10):1057-1068.
 98. Lovinsky-Desir S, Jung KH, Jezioro JR, et al. Physical activity, black carbon exposure, and DNA methylation in the FOXP3 promoter. *Clin Epigenetics*. 2017;9:65.
 99. Jung KH, Lovinsky-Desir S, Yan B, et al. Effect of personal exposure to black carbon on changes in allergic asthma gene methylation measured 5 days later in urban children: importance of allergic sensitization. *Clin Epigenetics*. 2017;9:61.
 100. Ziello C, Sparks TH, Estrella N, et al. Changes to airborne pollen counts across Europe. *PLoS ONE*. 2012;7(4):e34076.
 101. Suarez-Cervera M, Castells T, Vega-Maray A, et al. Effects of air pollution on cup a 3 allergen in Cupressus arizonica pollen grains. *Ann Allergy Asthma Immunol*. 2008;101(1):57-66.
 102. Donders TH, Hagemans K, Dekker SC, de Weger LA, de Klerk P, Wagner-Cremer F. Region-specific sensitivity of anemophilous pollen deposition to temperature and precipitation. *PLoS ONE*. 2014;9(8):e104774.
 103. Ziska LH, Gebhard DE, Frenz DA, Faulkner S, Singer BD, Straka JG. Cities as harbingers of climate change: common ragweed, urbanization, and public health. *J Allergy Clin Immunol*. 2003;111(2):290-295.
 104. Reinmuth-Selzle K, Kampf CJ, Lucas K, et al. Air pollution and climate change effects on allergies in the anthropocene: abundance, interaction, and modification of allergens and adjuvants. *Environ Sci Technol*. 2017;51(8):4119-4141.
 105. Garcia-Mozo H. Poaceae pollen as the leading aeroallergen worldwide: a review. *Allergy*. 2017;72(12):1849-1858.
 106. Carsin A, Romain T, Ranque S, et al. *Aspergillus fumigatus* in cystic fibrosis: an update on immune interactions and molecular diagnostics in allergic bronchopulmonary aspergillosis. *Allergy*. 2017;72(11):1632-1642.

107. Gilles-Stein S, Beck I, Chaker A, et al. Pollen derived low molecular compounds enhance the human allergen specific immune response in vivo. *Clin Exp Allergy*. 2016;46(10):1355-1365.
108. Beck I, Jochner S, Gilles S, et al. High environmental ozone levels lead to enhanced allergenicity of birch pollen. *PLoS ONE*. 2013;8(11):e80147.
109. Lang-Yona N, Shuster-Meiseles T, Mazar Y, Yarden O, Rudich Y. Impact of urban air pollution on the allergenicity of *Aspergillus fumigatus* conidia: outdoor exposure study supported by laboratory experiments. *Sci Total Environ*. 2016;541:365-371.
110. Cuinica LG, Abreu I, Esteves da Silva J. Effect of air pollutant NO₂ on *Betula pendula*, *Ostrya carpinifolia* and *Carpinus betulus* pollen fertility and human allergenicity. *Environ Pollut*. 2014;186:50-55.
111. Zhao F, Elkelish A, Durner J, et al. Common ragweed (*Ambrosia artemisiifolia* L.): allergenicity and molecular characterization of pollen after plant exposure to elevated NO₂. *Plant Cell Environ*. 2016;39(1):147-164.
112. Ackaert C, Kofler S, Horejs-Hoek J, et al. The impact of nitration on the structure and immunogenicity of the major birch pollen allergen Bet v 1.0101. *PLoS ONE*. 2014;9(8):e104520.
113. Baldacci S, Maio S, Cerrai S, et al. Allergy and asthma: effects of the exposure to particulate matter and biological allergens. *Respir Med*. 2015;109(9):1089-1104.
114. Annesi-Maesano I, Rouve S, Desqueyroux H, et al. Grass pollen counts, air pollution levels and allergic rhinitis severity. *Int Arch Allergy Immunol*. 2012;158(4):397-404.
115. Cakmak S, Dales RE, Coates F. Does air pollution increase the effect of aeroallergens on hospitalization for asthma? *J Allergy Clin Immunol*. 2012;129(1):228-231.
116. Guilbert A, Cox B, Bruffaerts N, et al. Relationships between aeroallergen levels and hospital admissions for asthma in the Brussels-Capital Region: a daily time series analysis. *Environ Health*. 2018;17(1):018-0378.
117. Seinfeld JH, Pandis SN. *Atmospheric Chemistry and Physics*, 2nd ed. Hoboken, NJ: John Wiley; 2006.
118. Council NR. America's Climate Choices: Panel on Advancing the Science of Climate Change; 2010. Report No.: ISBN 0-309-14588-0.
119. D'Amato G, Cecchi L, D'Amato M, Annesi-Maesano I. Climate change and respiratory diseases. *Eur Respir Rev*. 2014;23(132):161-169.
120. Galan C, Alcazar P, Oteros J, et al. Airborne pollen trends in the Iberian Peninsula. *Sci Total Environ*. 2016;550:53-59.
121. Storkey J, Stratonovitch P, Chapman DS, Vidotto F, Semenov MA. A process-based approach to predicting the effect of climate change on the distribution of an invasive allergenic plant in Europe. *PLoS ONE*. 2014;9(2):e88156.
122. Pfaar O, Bastl K, Berger U, et al. Defining pollen exposure times for clinical trials of allergen immunotherapy for pollen-induced rhinoconjunctivitis - an EAACI position paper. *Allergy*. 2017;72(5):713-722.
123. Ziska L, Knowlton K, Rogers C, et al. Recent warming by latitude associated with increased length of ragweed pollen season in central North America. *Proc Natl Acad Sci USA*. 2011;108(10):4248-4251.
124. Garcia-Mozo H, Oteros JA, Galan C. Impact of land cover changes and climate on the main airborne pollen types in Southern Spain. *Sci Total Environ*. 2016;549:221-228.
125. Lind T, Ekebom A, Alm Kubler K, Ostensson P, Bellander T, Lohmus M. Pollen season trends (1973-2013) in Stockholm Area, Sweden. *PLoS ONE*. 2016;11(11):e0166887.
126. Barbeau DN, Grimsley LF, White LE, El-Dahr JM, Lichtveld M. Mold exposure and health effects following hurricanes Katrina and Rita. *Annu Rev Public Health*. 2010;31:165-178. 161 p following 178.
127. Way DA, Montgomery RA. Photoperiod constraints on tree phenology, performance and migration in a warming world. *Plant Cell Environ*. 2015;38(9):1725-1736.
128. Beggs PJ. Impacts of climate change on aeroallergens: past and future. *Clin Exp Allergy*. 2004;34(10):1507-1513.
129. Shea KM, Truckner RT, Weber RW, Peden DB. Climate change and allergic disease. *J Allergy Clin Immunol*. 2008;122(3):443-453. quiz 454-445.
130. D'Amato G, Cecchi L. Effects of climate change on environmental factors in respiratory allergic diseases. *Clin Exp Allergy*. 2008;38(8):1264-1274.
131. Chapman DS, Haynes T, Beal S, Essl F, Bullock JM. Phenology predicts the native and invasive range limits of common ragweed. *Glob Chang Biol*. 2014;20(1):192-202.
132. Cunze S, Leiblein MC, Tackenberg O. Range expansion of *Ambrosia artemisiifolia* in Europe is promoted by climate change. *ISRN Ecol*. 2013. <http://dx.doi.org/10.1155/2013/610126>
133. Siroux V, Ballardini N, Soler M, et al. The asthma-rhinitis multimorbidity is associated with IgE polysensitization in adolescents and adults. *Allergy*. 2018;73(7):1447-1458.
134. Gao Z, Fu WY, Sun Y, et al. Artemisia pollen allergy in China: component-resolved diagnosis reveals allergic asthma patients have significant multiple allergen sensitization. *Allergy*. 2019;74(2):284-293.
135. Pinkerton KE, Rom WN, Akpinar-Elci M, et al. An official American Thoracic Society workshop report: climate change and human health. *Proc Am Thorac Soc*. 2012;9(1):3-8.
136. Takaro TK, Knowlton K, Balmes JR. Climate change and respiratory health: current evidence and knowledge gaps. *Expert Rev Respir Med*. 2013;7(4):349-361.
137. D'Amato G, Cecchi L, D'Amato M, Liccari G. Urban air pollution and climate change as environmental risk factors of respiratory allergy: an update. *J Investig Allergol Clin Immunol*. 2010;20(2):95-102. quiz following 102.
138. De Sario M, Katsouyanni K, Michelozzi P. Climate change, extreme weather events, air pollution and respiratory health in Europe. *Eur Respir J*. 2013;42(3):826-843.
139. D'Amato G, Holgate ST, Pawankar R, et al. Meteorological conditions, climate change, new emerging factors, and asthma and related allergic disorders. A statement of the World Allergy Organization. *World Allergy Organ J*. 2015;8(1):25.
140. Baccini M, Biggeri A, Accetta G, et al. Heat effects on mortality in 15 European cities. *Epidemiology*. 2008;19(5):711-719.
141. D'Ippoliti D, Michelozzi P, Marino C, et al. The impact of heat waves on mortality in 9 European cities: results from the EuroHEAT project. *Environ Health*. 2010;9:37.
142. Michelozzi P, Accetta G, De Sario M, et al. High temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir Crit Care Med*. 2009;179(5):383-389.
143. Lemmetyinen RE, Karjalainen JV, But A, et al. Higher mortality of adults with asthma: a 15-year follow-up of a population-based cohort. *Allergy*. 2018;73(7):1479-1488.
144. Analitis A, Michelozzi P, D'Ippoliti D, et al. Effects of heat waves on mortality: effect modification and confounding by air pollutants. *Epidemiology*. 2014;25(1):15-22.
145. North ML, Jones MJ, Maclsaac JL, et al. Blood and nasal epigenetics correlate with allergic rhinitis symptom development in the environmental exposure unit. *Allergy*. 2018;73(1):196-205.
146. Karatzas K, Katsifarakis N, Riga M, et al. New European Academy of Allergy and Clinical Immunology definition on pollen season mirrors symptom load for grass and birch pollen-induced allergic rhinitis. *Allergy*. 2018;73(9):1851-1859.
147. Karatzas K, Riga M, Berger U, Werchan M, Pfaar O, Bergmann KC. Computational validation of the recently proposed pollen season definition criteria. *Allergy*. 2018;73(1):5-7.
148. Fuertes E, Butland BK, Ross Anderson H, Carlsten C, Strachan DP, Brauer M. Childhood intermittent and persistent rhinitis

- prevalence and climate and vegetation: a global ecologic analysis. *Ann Allergy Asthma Immunol.* 2014;113(4):386-392.e389.
149. Surda P, Putala M, Siarnik P, Walker A, Bernic A, Fokkens W. Rhinitis and its impact on quality of life in swimmers. *Allergy.* 2018;73(5):1022-1031.
 150. Paterson RR, Lima N. Thermophilic fungi to dominate aflatoxigenic/mycotoxigenic fungi on food under global warming. *Int J Environ Res Public Health.* 2017;14(2):199.
 151. Altizer S, Ostfeld RS, Johnson PT, Kutz S, Harvell CD. Climate change and infectious diseases: from evidence to a predictive framework. *Science.* 2013;341(6145):514-519.
 152. Tormanen S, Lauhkonen E, Riikonen R, et al. Risk factors for asthma after infant bronchiolitis. *Allergy.* 2018;73(4):916-922.
 153. D'Amato G, Annesi-Maesano I, Cecchi L, D'Amato M. Latest news on relationship between thunderstorms and respiratory allergy, severe asthma, and deaths for asthma. *Allergy.* 2019;74(1):9-11.
 154. D'Amato G, Cecchi L, Annesi-Maesano I. A trans-disciplinary overview of case reports of thunderstorm-related asthma outbreaks and relapse. *Eur Respir Rev.* 2012;21(124):82-87.
 155. Cockcroft DW. Epidemic thunderstorm asthma. *Lancet Planet Health.* 2018;2(6):e236-e237.
 156. Hew M, Lee J, Susanto NH, et al. The 2016 Melbourne thunderstorm asthma epidemic: Risk factors for severe attacks requiring hospital admission. *Allergy.* 2019;74(1):122-130.
 157. Inspector General of Emergency Management, Victoria State Government. Review of Response to the Thunderstorm Asthma Event of 21-22 November 2016 Preliminary Report. [cited 2018 May 24]. https://www.igem.vic.gov.au/sites/default/files/embridge_cache/emshare/original/public/2017/07/d9/afdfc180a/Reviewofresponsetothunderstormasthmaeventof2122NovemberPreliminaryReport.pdf. Accessed July 1, 2018.
 158. Brunekreef B, Annesi-Maesano I, Ayres JG, et al. Ten principles for clean air. *Eur Respir J.* 2012;39(3):525-528.
 159. Brunekreef B, Kunzli N, Pekkanen J, et al. Clean air in Europe: beyond the horizon? *Eur Respir J.* 2015;45(1):7-10.
 160. Crossman-Barnes CJ, Peel A, Fong-Soe-Khioe R, Sach T, Wilson A, Barton G. Economic evidence for nonpharmacological asthma management interventions: a systematic review. *Allergy.* 2018;73(6):1182-1195.
 161. European Union's (EU) 2050 Climate strategies and target: Low carbon Economy. https://ec.europa.eu/clima/policies/strategies/2050_en. Accessed July 1, 2018.
 162. Lifting Europe's Dark Cloud: how cutting coal saves lives. https://www.wwf.de/fileadmin/fm-wwf/Publikationen-PDF/Lifting_Europe_s_Dark_Cloud.pdf. Accessed July 1, 2018.
 163. Annesi-Maesano I. The air of Europe: where are we going? *Eur Respir Rev.* 2017;26(146):170024.
 164. European Commission. Best Available Techniques (BAT) Reference Document for Large Combustion Plants. Industrial Emissions Directive 2010/75/EU (Integrated Pollution and Prevention. (Last checked: February 2019). http://eippcb.jrc.ec.europa.eu/reference/BREF/LCP/JRC_107769_LCPBref_2017.pdf. Accessed July 1, 2019.
 165. European Parliament and European Council. National Emission Ceilings (NEC) directive. Directive (EU) 2016/2284. (last checked February 2019) https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=urisrv:OJ.L_.2016.344.01.0001.01.ENG&toc=OJ.L:2016:344:TOC. Accessed July 1, 2019.
 166. Watts N, Amann M, Arnell N, et al. The 2018 report of the Lancet Countdown on health and climate change: shaping the health of nations for centuries to come. *Lancet.* 2018;392(10163):2479-2514.
 167. Lambert KA, Bowatte G, Tham R, et al. Residential greenness and allergic respiratory diseases in children and adolescents - A systematic review and meta-analysis. *Environ Res.* 2017;159:212-221.
 168. Cilluffo G, Ferrante G, Fasola S, et al. Associations of greenness, greyness and air pollution exposure with children's health: a cross-sectional study in Southern Italy. *Environ Health.* 2018;17(1):018-0430.
 169. Parmes E, Pesce G, Sabel CE, et al. Influence of residential land cover on childhood allergic and respiratory symptoms and diseases: Evidence from 9 European cohorts. *Environ Res.* 2019;108953. <https://doi.org/10.1016/j.envres.2019.108953>. [Epub ahead of print].
 170. Abramson SL. Reducing environmental allergic triggers: policy issues. *J Allergy Clin Immunol Pract.* 2018;6(1):32-35.
 171. Carinanos P, Alcazar P, Galan C, Dominguez E. Environmental behaviour of airborne Amaranthaceae pollen in the southern part of the Iberian Peninsula, and its role in future climate scenarios. *Sci Total Environ.* 2014;471:480-487.
 172. European Parliament Interest Group on Allergy and Asthma. <http://www.eaaci.org/outreach/eu-activities/european-parliament-interest-group-on-allergy-and-asthma.html>. Accessed July 1, 2018.
 173. EAACI, EFA and the European Parliament Interest Group on allergy and asthma launch a political call to action in Europe. http://www.callallergyasthma.eu/C2A_Press_Release_United_Action_for_Allergy_and_Asthma.pdf. Accessed July 1, 2018.
 174. Kubesch NJ, Therning Jorgensen J, Hoffmann B, et al. Effects of leisure-time and transport-related physical activities on the risk of incident and recurrent myocardial infarction and interaction with traffic-related air Pollution: a Cohort Study. *J Am Heart Assoc.* 2018;7(15). <https://doi.org/10.1161/JAHA.118.009554>
 175. Arbillaga-Etxarri A, Gimeno-Santos E, Barberan-Garcia A, et al. Long-term efficacy and effectiveness of a behavioural and community-based exercise intervention (Urban Training) to increase physical activity in patients with COPD. A randomised controlled trial. *Eur Respir J.* 2018;52(4):1800063.
 176. Erbas B, Jazayeri M, Lambert KA, et al. Outdoor pollen is a trigger of child and adolescent asthma emergency department presentations: A systematic review and meta-analysis. *Allergy.* 2018;73(8):1632-1641.
 177. Bousquet J, Caimmi DP, Bedbrook A, et al. Pilot study of mobile phone technology in allergic rhinitis in European countries: the MASK-rhinitis study. *Allergy.* 2017;72(6):857-865.
 178. Buters J, Schmidt-Weber C, Oteros J. Next-generation pollen monitoring and dissemination. *Allergy.* 2018;73(10):1944-1945.
 179. Heinrich J. Influence of indoor factors in dwellings on the development of childhood asthma. *Int J Hyg Environ Health.* 2011;214(1):1-25.
 180. Global Initiative for Asthma (GINA) 2019 Main Report. <https://ginasthma.org/gina-reports/2019>. Accessed July 1, 2019.
 181. Hellings PW, Fokkens WJ, Bachert C, et al. Positioning the principles of precision medicine in care pathways for allergic rhinitis and chronic rhinosinusitis - A EUFOREA-ARIA-EPOS-AIRWAYS ICP statement. *Allergy.* 2017;72(9):1297-1305.
 182. Bousquet J, Schunemann HJ, Samolinski B, et al. Allergic Rhinitis and its Impact on Asthma (ARIA): achievements in 10 years and future needs. *J Allergy Clin Immunol.* 2012;130(5):1049-1062.
 183. Uriarte SA, Sastre J. Clinical relevance of molecular diagnosis in pet allergy. *Allergy.* 2016;71(7):1066-1068.
 184. Lombardi E, Simoni M, La Grutta S, et al. Effects of pet exposure in the first year of life on respiratory and allergic symptoms in 7-yr-old children. The SIDRIA-2 study. *Pediatr Allergy Immunol.* 2010;21(2 Pt 1):268-276.

How to cite this article: Eguluz-Gracia I, Mathioudakis AG, Bartel S, et al. The need for clean air: The way air pollution and climate change affect allergic rhinitis and asthma. *Allergy.* 2020;75:2170-2184. <https://doi.org/10.1111/all.14177>